

**TREATS
4ha**



GRAZERS

EFFECTIVE AGAINST DAMAGE FROM

RABBITS

PIGEONS

DEER

GEESE

- ✓ **Effective for spot treatments and headland spray**
- ✓ **Safe to use on edible crops**
- ✓ **Spray from early leaf emergence**
- ✓ **Tank mix options**

1 litre

Grazers contains calcium and other ingredients:
calcium chloride solution - 10% calcium oxide, 7.2% w/w calcium
For tank mix compatibility please check our website

www.grazers.co.uk

Application rate

Basic rate: 250ml in 100-250 litres of water per ha. **Double rate:** 500ml in 100-250 litres of water per ha (for high infestations). **Knapsack rate:** 20ml in 20 litres of water. **Vines:** 3 litres in 1500 litres of water per ha. **Top fruit:** 2 litres in 1000 litres of water per ha.

Application timing

Apply to *dry foliage* (avoid frost and strong sunlight) at any stage of growth during the growing season (very effective on first true leaf), immediately the first signs of damage is evident, or preferably to areas at high potential risk.

GRAZERS is rainfast in one hour and can remain effective for up to 6 weeks, but in certain situations, where crops are grazed, mowed, irrigated or exposed to prolonged heavy rainfall after spraying, repeat applications may be required sooner.

GRAZERS is very safe to use and beneficial to the plant.

Arable (Cereals, Oil Seed Rape, Root Crops, Game Cover)

Grazers can be applied safely and effectively to any crop at risk from damage.

Cost effective treatments

Rabbits: a headland spray (min. 24m boom) is generally sufficient unless severe pest numbers.

Pigeons: treat all areas at risk (eg. under powerlines, below roosts) at early leaf emergence, and weaker areas during late winter to stimulate re growth and allow crop to recover.

Note: It is recommended to apply with a non-ionic surfactant, eg. Activator 90

Horticulture (Salad/Veg, Ornamental Crops) / Forage (Maize, Grass, Root Crops)

Grazers can be applied safely and effectively to any crop at risk from damage.

N.B. Modules, liners & transplants are extremely vulnerable.

Spray at least twice at double strength before planting/putting out, allow the leaf to dry between applications. Spray again 2-3 days later in the field.

Grazers is specifically designed to stimulate growth of the crop beyond the susceptible stages for grazing damage.

Note: It is recommended to apply with a non-ionic surfactant, eg. Activator 90

Vines & Top Fruit

Spray to wet the fruit thoroughly at or before first sign of damage approx. 2 weeks before harvest.

Note: addition of a non-ionic surfactant is recommended.

Tank mix compatibility

For up to date tank mix options check farmers and growers tab on www.grazers.co.uk

Grazers 10% Calcium Concentrate

WARNING Causes serious eye irritation.

Wear protective gloves/eye protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove Contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

Grazers, Gill Bank, Ousby, Penrith, CUMBRIA CA10 1QA, UK Tel: 01768 800555

