



3 active ingredients and their effects

- Florasulam This active ingredient is absorbed through both the leaves and the roots where it inhibits plant enzymes, stopping cell division and causing chlorosis and necrosis of the plant.
- Fluroxypyr This powerful ingredient goes in through the foliar part of the weed and de-regulates plant growth causing it to malform and twist.
- Clopyralid The third active ingredient, like fluroxypyr, is absorbed through both foliage and root, causing malformations and twisting.

Once inside the target plant, Praxys is rainfast within one hour, protected from being washed away by rain or irrigation.





Praxys is a premium selective, systemic post-emergence herbicide for use on managed amenity turf, including lawns and amenity grassland, for the control of daisy, dandelion, clover, buttercup, ribwort, plantain and other broad-leaved weeds.

Praxys combines three modern active ingredients and two different modes of action. Rapidly entering target weeds, Praxys travels through the whole plant to give complete weed destruction.

The Praxys low-down

- ✓ Lowest dosage rates
- ✓ Lowest active ingredient loading
- Lowest packaging waste costs
- Lowest water volume

Praxys is the most powerful and costeffective option in the market today. It has an extremely low dosage rate that gives a low environmental loading of active ingredient.

This means the lowest packaging waste, lowest storage and lowest carriage costs of any product.

Cost-effective dosage rates

Praxys offers flexible dosage rates according to the weed spectrum to be controlled. It should be applied as a 'Medium' spray as defined by the BCPC system. Maximum of one application per year.

| Sprayer type | Rate of Praxys | Water volume | Area treated | |
|---------------------|---|-----------------|-----------------|--|
| | 1 litre | 200 litres | 1 hectare | |
| | Controls Bird's-foot-trefoil*, Black Medick, Bristly Oxtongue, Common Cat's-ear, Creeping Cinquefoil, Sorrel, White Clover | | | |
| D | 1.5 litres | 200 litres | 1 hectare | |
| Power sprayer | Above plus Common Daisy, Common Dandelion, Greater Plantain, Common Mouse-ear | | | |
| | 2.0 litres | 200 litres | 1 hectare | |
| | Above plus Ribwort Plantain*, Self Heal, Creeping Buttercup, Spear Thistle, Slender Speedwell | | | |
| Knapsack spraver | 10 ml | | | |
| | 15 ml | 2 litres | 100 m² | |
| sprayer | | | | |

Product characteristics

| Formulation type: | Liquid (EC) |
|-------------------|---|
| Technology: | Selective herbicide |
| Pack size: | 2 litres |
| Pack coverage: | 10,000 - 20,000 m² |
| MAPP: | 19820 |
| PCS: | 03510 |
| Contains: | 144 g/L fluroxypyr-meptyl, 80 g/L clopyralid and 2.5 g/L florasulam |

Application information

| Maximum individual dose: | 2 L/ha |
|----------------------------|------------|
| Water volume: | 200 L/ha |
| Maximum no. of treatments: | 1 per year |

Long application window

Apply when weeds are in active growth normally from March to October when the soil is moist. Do not apply in periods of drought unless irrigation is applied.

Avoid mowing three days before and after spraying to ensure sufficient weed leaf surface is present and to allow uptake and movement of Praxys within the weed.

Extensive trials programme

Praxys was compared to three standard herbicides in a total of 29 efficacy trials in the UK, France and Italy.

Its high efficacy and consistent performance led to label approval giving operators a clean, powerful option for the control of a wide range of weeds.

Kills tough weeds with the 3 most powerful actives

> Slender Speedwell

> Spear Thistle

Creeping Buttercup

> White Clover

Dandelion

> Self Heal

Black Medick

Common Mouse-ear

> Ribwort Plantain

Daisy

Application notes

- The ideal timing for application is when the weeds are small and actively growing
- Broad-leaved weeds not present at application will not be controlled
- An interval of 4 weeks must elapse between application of Praxys and over-seeding turf
- Take extreme care to avoid drift onto crops and non-target plants, e.g. trees, shrubs and bedding outside the target area
- Do not use any plant material treated with Praxys for composting or mulching
- 🗙 Do not apply if turfgrass is wet
- Do not apply to turf, lawns or grass areas which are under stress
- Do not apply if night temperatures are low, if ground frost is imminent, or in periods of prolonged cold or dry weather

Always read the label and product information before use.

Areas of use

Managed amenity turf, including lawns and amenity grassland.

- Greens
- Tees
- Fairways
- Greens surrounds
- Greens collars & approaches
- Stadia pitches
- Sports pitches
- Outfields
- Racecourses & gallops
- Lawns
- Turf production
- Airfields
- Rough/semi rough
- Amenity grassland
- Cricket tables
- Lawn Tennis

Key:

- Ideal for purpose
- Suitable for purpose

Epsilon House West Road Ipswich Suffolk IP3 9FJ

United Kingdom

+44 1473 237 100

marketing.ukire@icl-group.com

9 0 in

www.icl-group.com

